

## Calibration of Geometric Distortion in the ACS Detectors

G. R. Meurer

*Department of Physics and Astronomy, The Johns Hopkins University, Baltimore, MD 21218*

D. Lindler

*Sigma Space Corporation, Lanham, MD 20706*

J. P. Blakeslee

*Department of Physics and Astronomy, The Johns Hopkins University, Baltimore, MD 21218*

C. Cox

*Space Telescope Science Institute, 3700 San Martin Drive, Baltimore, MD 21218*

A. R. Martel, H. D. Tran

*Department of Physics and Astronomy, The Johns Hopkins University, Baltimore, MD 21218*

R. J. Bouwens

*UCO/Lick Observatory, University of California, Santa Cruz, CA 95064*

H. C. Ford

*Department of Physics and Astronomy, The Johns Hopkins University, Baltimore, MD 21218*

M. Clampin, G. F. Hartig

*Space Telescope Science Institute, 3700 San Martin Drive, Baltimore, MD 21218*

M. Sirianni

*Department of Physics and Astronomy, The Johns Hopkins University, Baltimore, MD 21218*

G. de Marchi

*Space Telescope Science Institute, 3700 San Martin Drive, Baltimore, MD 21218*

**Abstract.** The off-axis location of the Advanced Camera for Surveys (ACS) is the chief (but not sole) cause of strong geometric distortion in all detectors: the Wide Field Camera (WFC), High Resolution Camera (HRC), and Solar Blind Camera (SBC). Dithered observations of rich star cluster fields are used to calibrate the distortion. We describe the observations obtained, the algorithms used to perform the calibrations and the accuracy achieved.

## 1. Introduction

Images from the *Hubble Space Telescope* (*HST*) Advanced Camera for Surveys (ACS) suffer from strong geometric distortion: the square pixels of its detectors project to trapezoids of varying area across the field of view. The tilted focal surface with respect to the chief ray is the primary source of distortion of all three ACS detectors. In addition, the *HST* Optical Telescope Assembly induces distortion as does the ACS M2 and IM2 mirrors (which are designed to remove *HST*'s spherical aberration). The SBC's optics include a photo-cathode and micro-channel plate which also induce distortion.

Here we describe our method of calibrating the geometric distortion using dithered observations of star clusters. The distortion solutions we derived are given in the IDC tables delivered in Nov 2002, and are currently implemented in the STScI CALACS pipeline. This paper is a more up to date summary of our results than that presented at the workshop. An expanded description of our procedure is given by Meurer (2002).

## 2. Method

### 2.1. Observations

The ACS SMOV geometric distortion campaign consisted of two *HST* observing programs: 9028 which targeted the core of 47 Tucanae (NGC 104) with the WFC and HRC, and 9027 which consisted of SBC observations of NGC 6681. Additional observations from programs 9011, 9018, 9019, 9024 and 9443 were used as additional sources of data, to check the results, and to constrain the absolute pointing of the telescope.

The CCD exposures of 47 Tucanae were designed to detect stars on the main sequence turn-off at  $m_B = 17.5$  in each frame. This allows for a high density of stars with relatively short exposures. The F475W filter (Sloan  $g'$ ) was used for the CCD observations so as to minimize the number of saturated red giant branch stars in the field. For the HRC two 60s exposures were taken at each pointing, while for the WFC which has a larger time overhead, only one such exposure was obtained per pointing. Simulated images made prior to launch, as well as archival WFPC2 images from Gilliland et al. (2000) were used to check that crowding would not be an issue. For calibrating the distortion in the SBC we used exposures of NGC 6681 (300s–450s) which was chosen for the relatively high density of UV emitters (hot horizontal branch stars). The pointing center was dithered around each star field. For the WFC and HRC pointings, the dither pattern was designed so that the offsets between all pairs of images adequately, and non-redundantly, samples all spatial scales from about 5 pixels to 3/4 the detector size. For the SBC pointings, a more regular pattern of offsets is used augmented by a series of 5 pixel offsets.

### 2.2. Distortion Model

The heart of the distortion model relates pixel position  $(x, y)$  to sky position using a polynomial transformation (Hack & Cox, 2000) given by:

$$x_c = \sum_{m=0}^k \sum_{n=0}^m a_{m,n} (x - x_r)^n (y - y_r)^{m-n}, \quad y_c = \sum_{m=0}^k \sum_{n=0}^m b_{m,n} (x - x_r)^n (y - y_r)^{m-n} \quad (1)$$

Here  $k$  is the order of the fit,  $x_r, y_r$  is the reference pixel, taken to be the center of each detector, or WFC chip, and  $x_c, y_c$  are undistorted image coordinates. The coefficients to the fits,  $a_{m,n}$  and  $b_{m,n}$ , are free parameters. For the WFC, an offset is applied to get the two CCD chips on the same coordinate system:

$$X' = x_c + \Delta x(\text{chip}\#), \quad Y' = y_c + \Delta y(\text{chip}\#). \quad (2)$$





Table 1. Summary of fit results

Camera	chip	pixel size [arcsec]	Filter	Pointings	$N$	rms(x) <sup>1</sup> [pixels]	rms(y) <sup>1</sup> [pixels]	Notes
WFC	1	0.05	F475W	25	142289	0.042	0.045	
WFC	2	0.05	F475W	25	103453	0.035	0.037	
WFC	1	0.05	F775W	10	31652	0.050	0.056	2
WFC	2	0.05	F775W	10	33834	0.041	0.048	2
HRC		0.025	F475W	20	77433	0.027	0.026	
HRC		0.025	F775W	13	31515	0.026	0.043	3
HRC		0.025	F220W	12	14715	0.112	0.108	3
SBC		0.03	F125LP	34	1561	0.109	0.094	

<sup>1</sup>This is the rms after iteratively clipping measurements with deviations greater than 5 times the rms.

<sup>2</sup>Coefficients held fixed to those found for WFC F475W.

<sup>3</sup>Coefficients held fixed to those found for HRC F475W.

and linear terms (scale, skewness). So we used the largest commanded offsets with a given guide star pair to set the linear terms. However, comparison of corrected coordinates to astrometric positions showed that residual skewness in the solution remained. Hence, as of November 2002, the IDC tables for WFC and SBC are based on data from multiple roll angles. The overall plate scale is set by the largest commanded offset. For the HRC, the linear scale is set by matching HRC and WFC coordinates, since the same field was used in the SMOV observations. The zeroth order terms (position of the ACS apertures in the *HST* V2, V3 frame) was determined from observations of an astrometric field.

### 3. Results

The distortion in all ACS detectors is highly non-linear as illustrated in Figure 1. We find that a quartic fit ( $k = 4$ ) is adequate for characterizing the distortion to an accuracy much better than our requirement of 0.2 pixels over the entire field of view. Table 1 summarizes the rms of the fits to the various datasets.

The WFC and HRC fits were all to F475W data as noted above. To check the wavelength dependence of the distortion we used data obtained with F775W (WFC and HRC) and F220W (HRC) from programs 9018 and 9019. We held the coefficients fixed and only fit the offsets in order to check whether a single distortion solution is sufficient for each detector. Table 2 shows that there is a marginal increase in the rms for the red data of the WFC, little or no increase in the fit rms for the red HRC data, but a significant increase in the rms using the UV data. An examination of the HRC F220W images reveals the most likely cause: the stellar PSF is elongated by 0.1". A similar elongation can also be seen in SBC PSFs. We attribute this to aberration in the optics of either the ACS M1 or M2 mirrors or the *HST* OTA (Hartig et al. 2002). The aberration amounts to 0.1 waves at 1600 Å, but is negligible relative to optical wavelengths, hence it is not apparent in optical HRC images. While it was expected that the same distortion solution would be applicable to all filters except the polarizers, recent work (by Tom Brown, STScI, and our team) has shown that at least one other optical filter (F814W) induces a significant plate scale change (factor of  $\sim 4 \times 10^{-5}$ ). In the long term, the IDC tables will be selected by filter in the STScI CALACS pipeline.

